PATIENT INSTRUCTIONS

Boston® IV (itafocon B)

Boston[®] II (itafocon A)

Silicone Acrylate Rigid Gas Permeable Contact Lenses For Daily Wear



CAUTION: Federal (USA) Law restricts this device to sale by or on the order of a licensed practitioner.

TABLE OF CONTENTS

Introduction

Wearing Restrictions and Indications

Contraindications

Warnings Precautions

Adverse Effects

Personal Cleanliness and Lens Handling

Preparing the Lens for Wearing

Handling the Lenses

Placing the Lens on the Eye

Centering the Lens

Removing the Lens

Caring for Your Lenses
(Cleaning, Rinsing, Disinfecting, Storage

and Rewetting/Lubricating)

Basic Instructions

Care for a Sticking (Nonmoving) Lens Lens Case Cleaning and Maintenance

Emergencies

Instructions for Monovision Wearer

Prescribed Wearing Schedule

Appointment Schedule

INTRODUCTION

Your eyecare practitioner has fit you with contact lenses known as Boston® IV or Boston® II. These lenses are manufactured from Boston IV and Boston II lens material. It is essential that you follow the recommended handling, cleaning and storage procedures. Failure to do so may eventually impair the performance of your lenses.

Boston® IV (itafocon B) and Boston® II (itafocon A) are Rigid Gas Permeable Contact Lens materials, composed of methyl-methacrylate-dimethylitaconate-siloxanyl copolymer. Boston IV lens materials are available in clear and blue. The blue Boston IV lens materials contain the color additive D & C Green No. 6. Boston II lens materials are available in clear, blue and green. The blue Boston II lens material contain the color additive D & C Green No. 6; the green Boston II lens material contain the color additive D & C Green No. 6 and C.I. Solvent Yellow No. 18.

WEARING RESTRICTIONS AND INDICATIONS

Boston IV and Boston II Contact Lenses are indicated for daily wear for the correction of refractive ametropia (myopia, hyperopia, astigmatism, presbyopia and keratoconus) in aphakic and not aphakic persons with nondiseased eyes. The lens may be disinfected using a chemical disinfection (not heat) system only.

Boston IV and Boston II contact lenses described in this booklet should be removed from your eyes for routine cleaning and disinfecting as prescribed by your eyecare practitioner. DO NOT WEAR YOUR THE BOSTON IV Or BOSTON II CONTACT LENSES WHILE SLEEPING.

CONTRAINDICATIONS (REASONS NOT TO USE)

DO NOT USE BOSTON IV or BOSTON II Contact Lenses when any of the following conditions exist:

- Acute or subacute inflammation or infection of the anterior chamber of the eye
- Any eye disease, injury, or abnormality that affects the cornea, conjunctiva, or eyelids
- Severe insufficiency of lacrimal secretion (dry eyes)
- Corneal hypoesthesia (reduced corneal sensitivity), if not aphakic
- Any systemic disease that may affect the eye or be exaggerated by wearing contact lenses
- Allergic reactions of ocular surfaces or adnexa that may be induced or exaggerated by wearing contact lenses or using contact lens solutions
- Allergy to any ingredient in a solution which is to be used to care for Boston IV or Boston II contact lenses
- Any active corneal infection (bacterial, fungal, or viral)
- · Red or irritated eyes

WARNINGS

- Problems with contact lenses and lens care products could result in serious injury to the eye. It is essential to follow your eyecare practitioner's directions and all labeling instructions for proper use of lenses and lens care products, including the lens case. Eye problems, including corneal ulcers, can develop rapidly and lead to loss of vision.
- Daily wear lenses are not indicated for overnight wear, and should not be worn while sleeping.
 Clinical studies have shown that the risk of serious adverse reactions is increased when these lenses are worn overnight.
- Studies have shown that contact lens wearers who are smokers have a higher incidence of adverse reactions than nonsmokers.
- If you experience eye discomfort, excessive tearing, vision changes, or redness of the eye, immediately remove lenses and promptly contact your evecare practitioner.

PRECAUTIONS

You should carefully adhere to the following care regimen and safety precautions:

 Before leaving the eyecare practitioner's office, you should be able to properly remove lenses or should have someone else available who can remove the lenses for you.

- You should remove your lenses immediately if your eyes become red or irritated.
- Different solutions often cannot be used together, and not all solutions are safe for use with all lenses. Use only recommended solutions.
 - Do not heat the wetting/soaking solution and lenses. Keep them away from extreme heat.
 - Always use fresh unexpired lens care solutions.
 - Always follow directions in the package insert for the use of contact lens solutions.
 - Use only a chemical (not heat) lens care system.
 Use of a heat (thermal) care system can warp the Boston® IV and Boston® II Contact Lenses.
 - Sterile unpreserved solutions, when used, should be discarded after the time specified in the labeling directions.
 - Do not use saliva or anything other than the recommended solutions for lubricating or wetting lenses.
 - Always keep the lenses completely immersed in the recommended storage solution when the lenses are not being worn (stored). If dry storage is desired to store the lenses for a longer period of time, they must first be cleaned, rinsed with water and carefully dried by blotting with a soft lint-free tissue prior to being placed in a clean, dry lens storage case. Ideally, these lenses should be cleaned and disinfected overnight prior to insertion.
- If the lens sticks (stops moving) on the eye, follow the recommended directions on Care for a sticking Lens. The lens should move freely on the eye for the continued health of the eye. If nonmovement of the lens continues, immediately consult your eyecare practitioner.
- Always wash and rinse hands before handling lenses. Do not get cosmetics, lotions, soaps, creams, deodorants, or sprays in the eyes or on the lenses. It is best to put on lenses before putting on makeup. Water-based cosmetics are less likely to damage lenses than oil-based products.
- Do not touch contact lenses with the fingers or hands if the hands are not free of foreign materials, as microscopic scratches on the lenses may occur, causing distorted vision and/or injury to the eye.
- Carefully follow the handling, insertion, removal, cleaning, disinfecting, storing and wearing instructions that follow for Boston IV and Boston II Contact Lenses and those provided by your eyecare practitioner.
- Never wear lenses beyond the period recommended by your eyecare practitioner.
- If aerosol products such as hair spray are used while wearing lenses, exercise caution and keep eyes closed until the spray has settled.
- Always handle lenses gently and avoid dropping them on hard surfaces.
- Avoid all harmful or irritating vapors and fumes while wearing lenses.
- Ask your eyecare practitioner about wearing lenses during water activities and other sports.
- Inform your health care practitioner (doctor) that you wear contact lenses.

- Never use tweezers or other tools to remove lenses from the lens case unless specifically indicated for that use. To remove the lens from the case, pour the solution containing the lens into the palm of your hand.
- . Do not touch the lens with fingernails.
- Always contact your eyecare practitioner before using any medicine in the eyes.
- Always inform your employer that you wear contact lenses. Some jobs may require use of eye protection equipment or may require that you not wear contact lenses.
- As with any contact lens, follow-up visits are necessary to assure the continuing health of your eyes. Follow your eyecare practitioner's instruction as to a recommended follow-up schedule.

ADVERSE EFFECTS

The following problems may occur:

- Eyes stinging, burning, itching (irritation), or other eye pain
- Comfort is less than when lens was first placed on the eye
- Feeling of something in the eye such as a foreign body, scratched area
- . Excessive watering (tearing) of the eyes
- · Unusual eye secretions
- · Redness of the eves
- Reduced sharpness of vision (poor visual acuity)
- · Blurred vision, rainbows, or halos around objects
- · Sensitivity to light (photophobia)
- Dry eyes

If you notice any of the above:

Immediately remove lenses.

If the discomfort or problem stops, then look closely at the lens. If the lens is in any way damaged, do not put the lens back on the eye. Place the lens in the storage case and contact your eyecare practitioner. If the lens has dirt, an eyelash, or other foreign body on it, or the problem stops and the lens appears undamaged, you should thoroughly clean, rinse, and disinfect the lenses; then reinsert them. After reinsertion, if the problem continues, immediately remove the lenses and consult your eyecare practitioner.

When any of the above problems occur, a serious condition such as infection, corneal ulcer, neovas-cularization, or initis may be present. You should keep the lens off the eye and seek immediate professional identification of the problem and prompt treatment to avoid serious eve damage.

PERSONAL CLEANLINESS FOR LENS HANDLING

1. Preparing the Lens for Wearing

- It is essential that you learn and use good hygienic methods in the care and handling of your new lenses. Cleanliness is the first and most important aspect of proper contact lens care. In particular, your hands should be clean and free of any foreign substances when you handle your lenses. The procedures are:
- Always wash your hands thoroughly with a mild soap, rinse completely, and dry with a lint-free towel before touching your lenses.

- Avoid the use of soaps containing cold cream, lotion, or oily cosmetics before handling your lenses, since these substances may come into contact with the lenses and interfere with successful wearing.
- Handle the lenses with your fingertips, and be careful to avoid contact with fingernails. It is helpful to keep your fingernails short and smooth

Start off correctly by getting into the habit of always using proper hygiene procedures so that they become automatic.

2 Handling the Lenses

- Develop the habit of always working with the same lens first to avoid mix-ups.
- Remove the lens from its storage case and examine it to be sure that it is moist, clean, clear, and free of any nicks or cracks.

3. Placing the Lens on the Eve

After thoroughly washing and rinsing your hands, and after proper cleaning and conditioning of the lens, follow these steps to insert the lens:

- Remove the lens from its storage compartment
- Rinse the lens with fresh conditioning solution, if desired.
- Inspect the lens to be sure that it is clean, uni-
- Rub several drops of conditioning solution over the lens surfaces.

formly wet and free of debris.

- Place the lens on the top of the index finger of your dominant hand. Place the middle finger of the same hand close to the lower lash and hold down the lower lid.
- Use the forefinger or middle finger of your other hand to lift the upper lid and then place the lens on the eye. It is not necessary to press the lens against the eye.
- Gently release the lids and blink. The lens will center automatically. Always verify its proper position by checking your vision immediately after insertion.
- Use the same technique or reverse the hand when inserting the other lens

There are other methods of lens placement. If the above method is difficult for you, your eyecare practitioner will provide you with an alternate method

Note: If after placement of the lens, your vision is blurred, check for the following:

- The lens is not centered on the eye (see "Centering the Lens." next in this booklet).
- If the lens is centered, remove the lens (see "Removing the Lens" section) and check for the following:
 - a. Cosmetics or oils on the lens. Clean, rinse, disinfect, and place on the eye again.
 - b. The lens is on the wrong eye.

If you find that your vision is still blurred after checking the above possibilities, remove both lenses and consult your eyecare practitioner.

4. Centering the Lens

Very rarely, a lens that is on the cornea will be displaced onto the white part of the eye during lens wear. This can also occur during placement and removal of the lenses if the correct techniques are not performed properly. To center a lens follow one of the procedures below

• Close your eyelids and gently massage the lens into place through the closed lids.

0

 Gently push the off-centered lens onto the cornea while the eye is open using finger pressure on the upper or lower lid next to the edge of the lens.

5. Removing the Lens

Before removing your lenses, it is recommended that you have the following items available:

1) A lens storage case.

2) Two Bottle Care System

Boston ADVANCE® Cleaner or Boston® Cleaner. AND Boston ADVANCE® Comfort Formula Conditioning Solution or Boston® Conditioning Solution.

OR

One Bottle Care System

Boston SIMPLUS® Multi-Action Solution (Removes Protein, Cleans, Disinfects, Conditions & Cushions)

A clean towel.

Always remove the same lens first.

- a. Wash, rinse, and dry your hands thoroughly.
- b. There are two suggested methods of lens removal:

TWO-FINGER METHOD

- 1) Place a towel under your eye to catch the lens.
- Place the tip of the forefinger of one hand on the middle of the upper lid margin and the forefinger of the other hand on the middle of the lower lid margin.
- Press the lid margin inward and then together. The lens should be wedged out of your eye onto your hand or towel.
- 4) The lens may come out but remain on your eyelid or hand or be decentered onto the white part of your eye. If the latter occurs, recenter the lens onto your cornea before repeating the removal procedure.

BLINK METHOD

Seat yourself at a table covered with a clean towel and lean over until you are looking down at the surface.

- 1) Place your index finger at the outer junction of your upper and lower lids, stretch the skin outward and slightly upward. (Do not allow your lid to slide over the lens.)
- 2) Blink briskly. The lens will be pinched by the pressure of your eyelids and the lens will pop out onto the clean surface of the towel, or you may catch the lens in the palm of your hand.
- Remove the other lens by following the same procedure.

d. Follow the required lens care procedures described under the heading, CARING FOR YOUR LENSES (CLEANING, RINSING, DISINFECT-ING, STORAGE AND REWETTING/LUBRICATING).

Note: If these methods for removing your lenses are difficult for you, your eyecare practitioner will provide you with an alternate method.

CARING FOR YOUR LENSES (CLEANING, RINSING, DISINFECTING, STORAGE AND REWETTING/LUBRICATING)

1. Basic Instructions

For continued safe and comfortable wearing of your lenses, it is important that you first Clean and rinse, then disinfect your lenses after each removal, using the care regimen recommended by your eyecare practitioner. Cleaning and rinsing are necessary to remove mucus, secretions, films, or deposits which may have accumulated during wear. The ideal time to clean your lenses is immediately after removing them. Disinfecting is necessary to destroy harmful germs.

You should adhere to the recommended care regimen. Failure to follow the regimen may result in development of serious ocular complications as discussed in the WARNINGS section above.

If you require only vision correction, but will not or cannot adhere to a recommended care regimen for your lenses, or are unable to place and remove lenses and do not have someone available to place and remove them for you, you should not attempt to wear contact lenses.

When you first get your lenses, be sure you can place the lenses on your eyes and remove them while you are in your eyecare practitioner's office. At that time you will be provided with a recommended cleaning and disinfection regimen and instructions and warnings for lens care, handling, cleaning, and disinfection. Your eyecare practitioner should instruct you about appropriate and adequate procedures and products for your use, and provide you with a copy of the Patient Instructions for Boston® IV and Roston® II Contact lenses.

For safe contact lens wear, you should know and always practice your lens care routine:

- Always wash, rinse, and dry hands before handling contact lenses.
- Always use fresh unexpired lens care solutions.
- Use the recommended system of lens care (chemical not heat) and carefully follow instructions on solution labeling. Different solutions often cannot be used together, and not all solution are safe for use with all lenses. Do not alternate or mix lens care systems unless indicated on solution labeling, or if advised by your eyecare practitioner.
- Always remove, clean, rinse and disinfect your lenses according to the schedule prescribed by your eyecare practitioner. The use of an enzyme or any cleaning solution does not substitute for disinfection.
- Do not use saliva or anything other than the recommended solutions for lubricating or rewetting your lenses. Do not put lenses in your mouth.

The lens care products listed below are recommended by Bausch & Lomb for use with either your Boston IV or Boston IV Contact Lenses. Your eyecare practitioner may recommend alternate products that are appropriate for you to use with your Boston IV or Boston II Contact Lenses.

LENS CARE TABLE		
Product Purpose	Lens Care System Chemical (Not Heat)	
Clean	Boston ADVANCE® Cleaner or Boston® Cleaner	
Disinfect	Boston ADVANCE® Comfort Formula Conditioning Solution or Boston® Conditioning Solution	
Store	Boston ADVANCE Comfort Formula Conditioning Solution or Boston Conditioning Solution	
Multi-Action (Clean, Condition, Disinfect, Rinse and Cushion)	Boston SIMPLUS® Multi-Action Solution	
Lubricate/Rewet	Boston® Rewetting Drops	
Weekly Enzymatic Cleaner	Boston® ONE STEP Liquid Enzymatic Cleaner	

- Note: Some solutions may have more than one function, which will be indicated on the label. Read the label on the solution bottle, and follow instructions.
- Clean one lens first (always the same lens first to avoid mix-ups) and rinse the lens thoroughly as recommended by your eyecare practitioner to remove the cleaning solution, mucus, and film from the lens surface. Follow the instructions provided in the cleaning solution labeling. Put that lens into the correct chamber of the lens storage case. Then repeat the procedure for the second lens.
- After cleaning, disinfect lenses using the above recommended system by your eyecare practitioner and/or the manufacturer. Follow the instructions provided in the disinfection solution labeling.
- To store lenses, disinfect and leave them in the closed/unopened case until ready to wear. If lenses are not to be used immediately following disinfection, you should consult the Package Insert or your eyecare practitioner for information on storage of your lenses.
- Always keep your lenses completely immersed in a recommended disinfecting/conditioning solution when the lenses are not being worn.
 If you discontinue wearing your lenses, but plan to begin wearing them again after a few weeks, ask your eyecare practitioner for a recommendation on how to store your lenses.
- Boston IV and Boston II Contact Lenses cannot be heat (thermally) disinfected.
- After removing your lenses from the lens case, empty and rinse the lens storage case with solution(s) recommended by the lens case manufacturer or the eyecare practitioner; then allow the lens case to air dry. When the case is used again, refill it with fresh storage solution.

Lens cases should be replaced at regular intervals as recommended by the lens case manufacturer or your evecare practitioner.

- Your eyecare practitioner may recommend a lubricating/rewetting solution for your use. Lubricating/Rewetting solutions can be used to wet (lubricate) your lenses while you are wearing them to make them more comfortable.
- Your eyecare practitioner may recommend a weekly enzymatic cleaner which can be used to effectively remove protein deposits from your Boston® IV or Boston® II Rigid Gas Permeable Contact Lenses.

2. Care for a Sticking (Nonmoving) Lens

If the lens sticks (stops moving/cannot be removed), apply one to three drops of a recommended lubricating or rewetting solution directly to your eye and wait until the lens begins to move freely on the eye before removing it. If nonmovement of the lens continues after 5 minutes, you should immediately consult your eyecare practitioner.

3. Lens Case Cleaning and Maintenance

Contact lens cases can be a source of bacterial growth. Lens cases should be emptied, cleaned, rinsed with solutions recommended by the lens case manufacturer or the eyecare practitioner, and allowed to air dry after each use. Lens cases should be replaced at regular intervals as recommended by the lens case manufacturer or the eyecare practitioner.

4. Emergencies

If chemicals of any kind (household products, gardening solutions, laboratory chemicals, etc.) are splashed into your eyes, you should: FLUSH YOUR EYES IMMEDIATELY WITH TAP WATER, THEN REMOVE YOUR LENSES PROMPTLY, IF POSSIBLE, AND IMMEDIATELY CONTACT YOUR EYECARE PRACTITIONER OR VISIT A HOSPITAL

EMERGENCY ROOM WITHOUT DELAY. INSTRUCTIONS FOR THE MONOVISION WEARER

- . You should be aware that, as with any type of lens correction, there are advantages and compromises to monovision contact lens therapy. The benefit of clear near vision in all gazes that is available with monovision may be accompanied by a vision compromise that may reduce your visual acuity and depth perception for distance and near tasks. Some patients have experienced difficulty adapting to monovision. Symptoms, such as mild blurred vision, dizziness, headaches and a feeling of slight imbalance, may last for a brief minute or for several weeks as adaptation takes place. The longer these symptoms persist, the poorer is your prognosis for successful adaptation. You should avoid visually demanding situations during the initial adaptation period. It is recommended that you first wear these contact lenses in familiar situations which are not visually demanding. For example, be a passenger rather than a driver of an automobile during the first few days of lens wear. It is recommended that you only drive with monovision correction if you pass your state drivers' license requirements with monovision correction.
- Some monovision lens wearers will never be fully comfortable functioning under low levels of illumination, such as driving at night. If this happens, discuss with your eyecare practitioner whether you should have additional contact lenses pre-

- scribed so that both eyes are corrected for distance when sharp distance vision is required.
- If you require very sharp near vision during prolonged close work, you may want to discuss with your eyecare practitioner having additional contact lenses prescribed so that both eyes are corrected for near when sharp near vision is required.
- Some monovision lens wearers require supplemental spectacles to wear over the monovision correction to provide the clearest vision for critical tasks. You should discuss this with your eyecare practitioner.
- It is important that you follow your eyecare practitioner's suggestions for adaptation to monovision contact lens therapy. You should discuss any concerns that you may have during and after the adaptation period.
- The decision to be fit with a monovision correction is most appropriately left to the eyecare practitioner in conjunction with the patient after carefully considering and discussing your needs.

CONSIDERATIONS FOR BIFOCAL LENSES

Patients who are considering bifocal contact lenses should be highly motivated and must be informed of the benefits as well as the problems that may be encountered while adapting to bifocal contact lens wear.

Your eyecare practitioner may discuss the following with you:

1. Adaptation

Both bifocal spectacle and bifocal contact lens wearers need to learn to adapt to proper head positioning. The bifocal patient must position the head upright while rotating the eyes downward to read. Once the bifocal patient has adapted, proper positioning becomes effortless.

2. Driving at Night

Bifocal contact lens wearers should experience night vision before actually driving while wearing their lenses.

3. Flare at Night

Bifocal contact lenses wearers may experience flare at night. This may occur with certain lens designs. With time, bifocal contact lens wearers adapt to this situation.

4. Visual Expectation

Bifocal contact lens wearers may experience visual acuities less than could be achieved with bifocal spectacles.

SAMPLE OF WEARING AND APPOINTMENT SCHEDULES

Wearing Schedule

DAY	WEARING TIME (Hours)*
1	4 to 8 Hours
2	6 to 10 Hours
3	8 to 14 Hours
4	10 to 15 Hours
5	12 to All Waking Hours
6 and after	All Waking Hours
*if the lenses	continue to be well-tolerated.

ii ale lelibes contante to be well tolerated

<u>WARNING</u>: Boston IV and Boston II Contact Lenses are **NOT** intended for overnight (extended) wear.

APPOINTMENT SCHEDULE				
Minimum number	of hours lenses to be wo	rn at time of appointment:_		
Your appointment	s are on:			
Month	Year	Time	Date	
				_
PATIENT/EYECAR	E PRACTITIONER INFORM	IATION:		
Eyecare Practitione	er Information			
Practitioner Name				
Practice Name:				_
Practitioner Addre	SS:			
B	N			
Practitioner Phone	Number:			_
Perommended Le	ns Care Penimen			
Recommended Ec	is care regimen.			
Cleaning Solution	on:			
Conditioning So	olution:			
Rewetting Solu	tion:			
Weekly Enzyma	itic Cleaner:			

IMPORTANT: In the event that you experience any difficulty wearing your lenses or you do not understand the instructions given you, DO NOT WAIT for your next appointment. TELEPHONE YOUR EYECARE PRACTITIONER IMMEDIATELY.

© Bausch & Lomb Incorporated 1400 North Goodman Street Rochester, NY 14609 1-800-333-4730

® denotes registered trademarks of Bausch & Lomb Incorporated.

Print Date: 09/07 LB9135/06

PACKAGE INSERT

Boston® IV (itafocon B) Boston® II (itafocon A)

Spherical & Aspherical Contact
Lenses for Myopia & Hyperopia,
Keratoconus Contact Lenses,
Bifocal Contact Lenses for
Presbyopia and Toric Lenses
to Correct Astigmatism in
Not aphakic and Aphakic Persons

Silicone Acrylate Rigid Gas Permeable Contact Lenses For Daily Wear

IMPORTANT:

Please read carefully and keep this information for future use. This package insert is intended for the eyecare practitioner, but should be made available to patients upon request. The eyecare practitioner should provide the patient with the patient instructions that pertain to the patient's prescribed lens.

CAUTION:

Federal (USA) Law restricts this device to sale by or on the order of a licensed practitioner.

DESCRIPTION

Boston® IV (itafocon B) and Boston® II (itafocon A) are Rigid Gas Permeable Contact Lens materials, composed of methyl-methacrylate-dimethylita-conate-siloxanyl copolymer. Boston IV lens materials are available in clear and blue. The blue Boston IV lens materials contain the color additive D & C Green No. 6. Boston II lens materials are available in clear, blue and green. The blue Boston II lens material contain the color additive D & C Green No. 6; the green Boston II lens material contain the color additive D & C Green No. 6. and C.I. Solvent Yellow No. 18.

Boston IV & Boston II Contact Lenses are hemispherical shell of the following dimensions:

Spherical Lens Design		
Power Range	-20.00D to +20.00D	
	in 0.25D increments	
Diameter	7.0 mm to 11.5 mm	
Base Curve Range	5.00 mm to 9.00 mm	
	in .01 mm increments	
Aspherical Lens Designs		

(Some of these designs are patented; manufacture of these lenses in Boston IV (litafocon B) and/or Boston II (litafocon A) materials are authorized to licensed labs only)

Power Range -20.00D to +20.00D in 0.25D increments
Diameter 7.0 mm to 11.5 mm
Base Curve Range 6.00 mm to 9.20 mm in 0.01 mm increments

Bifocal Lens Designs

(Some of these designs are patented; manufacture of these lenses in Boston IV (itafocon B) and/or Boston II (itafocon A) materials are authorized to licensed labs only)

Power Range	-20.00D to +20.00D
	in 0.25D increments
Diameter	8.5 mm to 11.5 mm
Base Curve Range	6.30 mm to 9.50 mm
	in 0.01 mm increments
Segment Heights	-2.00 mm to +1.00 mm
	in 0.5 mm increments
Add Powers	+1.00D to +3.75D
	in 0.5D increments
Prism Ballast	0.5 to 3.5 prism diopter
	in 0.5D increments

Toric Lens Designs

Power Range	-20.00D to +20.00D
	in 0.25D increments
Diameter	7.0 mm to 11.5 mm
Base Curve Range	6.80 mm to 9.50 mm
	in 0.01 mm increments
Toricity	Up to 9.00 Diopters
Keratoconus Lens Designs	

Keratocorius	Lens Designs
Range	-20.00D to +20.00D

	in 0.25D increments
Diameter	7.0 mm to 11.5 mm
Base Curve Range	4.00 mm to 9.00 mm
	in 0.01 mm increments
Base Optic Zone	5.00 mm to 9.00 mm
	in 0.01 mm increments

Power

Boston IV

I III SIGAL I KOI LKIILS	
Hardness (Rockwell R)	117
Specific Gravity	1.10
Refractive Index	1.469
Light Transmittance (Above 500 nm)	>90%
Water Content	<1%
Oxygen Permeability	26* (18**)
Wetting Angle	17°

Boston II

PHYSICAL PROPERTIES

DHYSICAL DEODEDTIES

Hardness (Rockwell R)	119
Specific Gravity	1.13
Refractive Index	1.471
Light Transmittance (Above 500 nm)	>90%
Water Content	<1%
Oxygen Permeability	18* (11**
Wetting Angle	20°
*gas to gas method	

^{**}polarographic method (ISO/Fatt)

ACTIONS

Boston IV and Boston II Contact Lenses when placed on the cornea act as a refracting medium to focus light rays on the retina. The toric lens provides a more even surface over the uneven astigmatic cornea and thus helps to focus light rays on the retina.

INDICATIONS (USES)

These lenses are indicated for daily wear. These lenses are intended to correction of visual acuity in non-aphakic persons with non-diseased eyes that are myopic, hyperopic or presbyopic, or have keratoconus and may exhibit up to 4.00D of corneal astigmatism that does not interfere with visual acuity. The lenses may be disinfected using a chemical disinfection system only.

CONTRAINDICATIONS (REASONS NOT TO USE)

DO NOT USE either Boston IV or Boston II Contact Lenses when any of the following conditions exist:

- Acute or subacute inflammation of the anterior chamber of the eye
 Any eye disease, injury, or abnormality, other
- Any eye disease, injury, or abnormality, other than Keratoconus, that affects the cornea, conjunctiva, or evelids
- Severe insufficiency of lacrimal secretion (dry eyes)
- Corneal hypoesthesia (reduced corneal sensitivity), if not aphakic
- Any systemic disease that may affect the eye or be exaggerated by wearing contact lenses
- Allergic reactions of ocular surfaces or adnexa that may be induced or exaggerated by wearing contact lenses or using contact lens solutions
- Allergy to any ingredient in a solution which is to be used to care for Boston IV or Boston II contact lens materials.
- Any active corneal infection (bacterial, fungal, or viral)
- · Red or irritated eyes

WARNINGS

Patients should be advised of the following warnings pertaining to contact lens wear:

- Problems with contact lenses and lens care products could result in serious injury to the eye. It is essential that patients follow their eyecare practitioner's directions and all labeling instructions for proper use of lenses and lens care products, including the lens case. Eye problems, including corneal ulcers, can develop rapidly and lead to loss of vision.
- Daily wear lenses are not indicated for overnight wear, and patients should be instructed not to wear lenses while sleeping. Clinical studies have shown that the risk of serious adverse reactions is increased when these daily wear lenses are worn overnight.
- Studies have shown that contact lens wearers who are smokers have a higher incidence of adverse reactions than possmokers.
- If a patient experiences eye discomfort, excessive tearing, vision changes, or redness of the eye, the patient should be instructed to immediately remove lenses and promptly contact his or her eyecare practitioner.

PRECAUTIONS

Practitioner Note: Boston® IV and Boston® II contact lenses are not sterile when shipped from the Authorized Boston® Manufacturer. Prior to dispensing, clean and disinfect the lens(es) according to the appropriate lens care regimen.

Special Precautions for Eyecare Practitioners:

- Due to the small number of patients enrolled in clinical investigation of lenses, all refractive powers, design configurations, or lens parameters available in the lens material are not evaluated in significant numbers. Consequently, when selecting an appropriate lens design and parameters, the eyecare practitioner should consider all characteristics of the lens that can affect lens performance and ocular health, including oxygen permeability, wettability, central and peripheral thickness, and optic zone diameter.
- The potential impact of these factors on the patient's ocular health should be carefully weighed against the patient's need for refractive correction; therefore, the continuing ocular health of the patient and lens performance on the eye should be carefully monitored by the prescribing eyecare practitioner.
- Patients who wear contact lenses to correct presbyopia may not achieve the best corrected visual acuity for either far or near vision. Visual requirements vary with the individual and should be considered when selecting the most appropriate type of lens for each patient.
- Aphakic patients should not be fitted with either Boston IV or Boston II Contact Lenses until the determination is made that the eye has healed completely.
- Before leaving the eyecare practitioner's office, the patient should be able to properly remove lenses or should have someone else available who can remove the lenses for him or her.

 Eyecare practitioners should instruct the patient to remove the lenses immediately if the eye becomes red or irritated.

Eyecare practitioners should carefully instruct patients about the following care regimen and safety precautions:

- Different solutions often cannot be used together, and not all solutions are safe for use with all lenses. Use only recommended solutions.
 - Do not heat the conditioning/storage solution and/or lenses. Keep them away from extreme heat.
 - Always use fresh unexpired lens care solutions.
 - Always follow directions in the package inserts for the use of contact lens solutions.
 - Use only a chemical (not heat) lens care system. Use of a heat (thermal) care system can warp either the Boston IV or Boston II Contact Lenses.
 - Do not use saliva or anything other than the recommended solutions for lubricating or wetting lenses.
 - Sterile unpreserved solutions, when used, should be discarded after the time specified in the labeling directions.
 - Always keep the lenses completely immersed in the recommended storage solution when the lenses are not being worn (stored). If dry storage is desired to store the lenses for a longer period of time, they must first be deaned, rinsed with water and carefully dried by blotting with a soft lint-free tissue prior to being placed in a clean, dry lens storage case. Ideally, these lenses should be cleaned and disinfected prior to insertion.
- If the lens sticks (stops moving) on the eye, follow the recommended directions on Care for a sticking lens. The lens should move freely on the eye for the continued health of the eye. If nonmovement of the lens continues, the patient should be instructed to immediately consult his or her eyecare practitioner.
- Always wash and rinse hands before handling lenses. Do not get cosmetics, lotions, soaps, creams, deodorants, or sprays in the eyes or on the lenses. It is best to put on lenses before putting on makeup. Water-based cosmetics are less likely to damage lenses than oil-based products.
- Do not touch contact lenses with the fingers or hands if the hands are not free of foreign materials, as microscopic scratches on the lenses may occur, causing distorted vision and/or injury to the eye.
- Carefully follow the handling, insertion, removal, cleaning, disinfecting, storing and wearing instructions in the Patient Instructions for Boston IV and Boston II Contact Lenses and in those prescribed by the eyecare practitioner.
- Never wear lenses beyond the period recommended by the eyecare practitioner.

- If aerosol products such as hair spray are used while wearing lenses, exercise caution and keep eyes closed until the spray has settled.
- Always handle lenses gently and avoid dropping them on hard surfaces.
- Avoid all harmful or irritating vapors and fumes while wearing lenses.
- Ask your eyecare practitioner about wearing lenses during water activities and other sports.
- Instruct patient to inform his or her health care practitioner (doctor) that the patient wears contact lenses.
- Never use tweezers or other tools to remove lenses from the lens case unless specifically indicated for that use. Pour the lens into the hand.
- . Do not touch the lens with fingernails.
- Instruct the patient to contact his or her eyecare practitioner before using any medicine in the eyes.
- Instruct the patient to inform his or her employer that he or she wears contact lenses. Some jobs may require use of eye protection equipment or may require that the patient not wear contact lenses.
- As with any contact lens, follow-up visits are necessary to assure the continuing health of the patient's eyes. The patient should be instructed as to a recommended follow-up schedule.

ADVERSE EFFECTS

The patient should be informed that the following problems may occur:

- Eyes stinging, burning, itching (irritation), or other eye pain
- Comfort is less than when lens was first placed on the eve
- Feeling of something in the eye such as a foreign body, scratched area
- Excessive watering (tearing) of the eyes
- Unusual eye secretions
- Redness of the eyes
- Reduced sharpness of vision (poor visual acuity)
- Blurred vision, rainbows, or halos around objects
- Sensitivity to light (photophobia)
- Dry eyes

If the patient notices any of the above, he or she should be instructed to:

Immediately remove lenses.

• If the discomfort or problem stops, then look closely at the lens. If the lens is in any way damaged, do not put the lens back on the eye. Place the lens in the storage case and contact the eyecare practitioner. If the lens has dirt, an eyelash, or other foreign body on it, or the problem stops and the lens appears undamaged, the patient should thoroughly clean, rinse, and disinfect the lenses; then reinsert them. After reinsertion, if the problem continues, immediately remove the lenses and consult the evecare practitioner.

When any of the above problems occur, a serious condition such as infection, comeal ulcer, neovas-cularization, or iritis may be present. The patient should be instructed to **keep the lens off the eye and seek immediate** professional identification of the problem and prompt treatment to avoid serious eye damage.

FITTING

Conventional methods of fitting contact lenses apply to Boston® IV (itafocon B) and Boston® (itafocon A). Contact Lenses. For a detailed description of the fitting techniques, refer to Boston IV/Boston II Professional Fitting and Information Guide, copies of Which are available from:

Practitioner Marketing Representative Boston Products Group of Bausch & Lomb 1400 North Goodman Street Rochester, NY 14609 1-800-275-1241

Professional Fitting Guides are also available through your Authorized Boston® Manufacturer.

WEARING SCHEDULE

The wearing and replacement schedules should be determined by the eyecare practitioner. Patients tend to overwear the lenses initially.

The eyecare practitioner should emphasize the importance of adhering to the initial maximum wearing schedule. Regular checkups, as determined by the eyecare practitioner, are also extremely important.

Boston IV and Boston II Contact Lenses are indicated for daily wear. The suggested wearing time for these lenses is:

DAY	WEARING TIME (Hours)*	
1	4 to 8 hours	
2	6 to 10 hours	
3	8 to 14 hours	
4	10 to 15 hours	
5	12 to All Waking Hours	
6 and after	All Waking Hours	
*if the lenses continue to be well-tolerated.		

<u>WARNING</u>: Boston IV and Boston II Contact Lenses are **NOT** intended for overnight (extended) wear.

LENS CARE DIRECTIONS

Eyecare practitioners should review with the patient lens care directions, including both basic lens care information and specific instructions on the lens care regimen recommended for the patient:

General Lens Care (First Clean and Rinse, Then Disinfect Lenses) Basic Instructions:

Always wash, rinse, and dry hands before handling contact lenses.

- · Always use fresh unexpired lens care solutions.
- Use the recommended system of lens care, chemical (not heat) and carefully follow instructions on solution labeling. Different solutions often cannot be used together, and not all solutions are safe for use with all lenses. Do not alternate or mix lens care systems unless

alternate or mix lens care systems unless indicated on solution labeling, or if advised by the eyecare practitioner.

- Do not use saliva or anything other than the recommended solutions for lubricating or rewetting lenses. Do not put lenses in the mouth.
- Lenses should be cleaned, rinsed, and disinfected each time they are removed. Cleaning and rinsing are necessary to remove mucus and film from the lens surface. Disinfecting is necessary to destroy harmful germs.
- Always remove, clean, rinse, and disinfect lenses according to the schedule prescribed by the eyecare practitioner. The use of an enzyme or a cleaning solution does not substitute for disinfection.

The lens care products listed below are recommended by Bausch & Lomb for use with either Boston IV or Boston II Contact Lenses. Eyecare practitioners may recommend alternate products that are appropriate for the patient's use with his or her lensles).

LENS CARE TABLE

LENS CARE TABLE	
Product Purpose	Lens Care System Chemical (Not Heat)
Clean	Boston ADVANCE® Cleaner or Boston® Cleaner
Disinfect	Boston ADVANCE® Comfort Formula Conditioning Solution or Boston® Conditioning Solution
Store	Boston ADVANCE® Comfort Formula Conditioning Solution or Boston® Conditioning Solution
Multi-Action (Clean, Condition, Disinfect, Rinse and Cushion)	Boston SIMPLUS® Multi-Action Solution
Lubricate/Rewet	Boston® Rewetting Drops
Weekly Enzymatic	Boston® ONE STEP Liquid Enzymatic Cleaner

 Note: Some solutions may have more than one function, which will be indicated on the label.
 Read the label on the solution bottle, and follow instructions.

Cleaner

- Clean one lens first (always the same lens first to avoid mix-ups), rinse the lens thoroughly as directed by your eyecare practitioner to remove the cleaning solution, mucus, and film from the lens surface, and put that lens into the correct chamber of the lens storage case.
 Then repeat the procedure for the second lens.
- After cleaning, disinfect lenses using the system recommended by the manufacturer and/or the eyecare practitioner. Follow the instructions provided in the disinfecting solution packaging.
- To store lenses, disinfect and leave them in the closed/unopened case until ready to wear. If lenses are not to be used immediately following disinfection, the patient should be instructed to consult the package insert or the eyecare practitioner for information on storage of lenses.
- After removing the lenses from the lens case, empty and rinse the lens storage case with solution as recommended by the lens case manufacturer; then allow the lens case to air dry. When the case is used again, refill it with storage solution. Replace lens case at regular intervals as recommended by the lens case manufacturer or your eyecare practitioner.
- Eyecare practitioners may recommend a lubricating/rewetting solution which can be used to wet (lubricate) lenses while they are being worn to make them more comfortable.
- Eyecare practitioners may recommend a weekly enzymatic cleaner which can be used to effectively remove protein deposits from Boston® IV or Boston® II Contact Lenses.
- Boston IV and Boston II Contact Lenses cannot be heat (thermally) disinfected.

LENS CASE CLEANING AND MAINTENANCE

Contact lens cases can be a source of bacterial growth. Lens cases should be emptied, cleaned, rinsed with solutions recommended by the lens case manufacturer or the eyecare practitioner, and allowed to air dry. Lens cases should be replaced at regular intervals as recommended by the lens case manufacturer or the eyecare practitioner.

CARE FOR A STICKING (NONMOVING) LENS

If the lens sticks (stops moving/cannot be removed), the patient should be instructed to apply one to three drops of a recommended lubricating or rewetting solution directly to the eye and wait until the lens begins to move freely on the eye before removing it. If nonmovement of the lens continues after 5 minutes, the patient should immediately consult the eyecare practitioner.

EMERGENCIES

The patients should be informed that if chemicals of any kind (household products, gardening solutions, laboratory chemicals, etc.) are splashed into the eyes, the patient should: FLUSH EYES IMMEDIATELY WITH TAP WATER, THEN REMOVE LENSES PROMPTLY, IF POSSIBLE, AND IMMEDIATELY CONTACT THE EYECARE PRACTITIONER OR VISIT A HOSPITAL EMERGENCY ROOM WITHOUT DELAY.

HOW SUPPLIED

Each lens is supplied (nonsterile) in a plastic lens storage case. The case is labeled with the base curve, diopter power, diameter, center thickness, color, UV-absorber (if present) and lot number. Additional parameters of add power, segment height, prism ballast and truncation may be included for bifocal lenses.

REPORTING OF ADVERSE REACTIONS

All serious adverse reactions observed in patients wearing either Boston® IV or Boston® II Contact Lenses or adverse experiences with the lenses should be reported to:

Consumer Affairs Bausch & Lomb 1400 North Goodman Street Rochester, NY 14609 1-800-333-4730

© Bausch & Lomb Incorporated 1400 North Goodman Street Rochester, NY 14609 1-800-333-4730

® denotes registered trademarks of Bausch & Lomb Incorporated

Print Date: 09/07 LB9135/06

THIS COMPONENT PRINTED TWO SIDES

SPECIAL INSTRUCTIONS		BAUSCH & LOMB Vision care	APPROVALS / DATE PKG/GRPH
DIE LINE DOES NOT PRINT Printed bar codes must be vei	COATING PER SPECIFICATION RIFIED READABLE BY PRINTING VENDOR	PACKAGING/GRAPHIC ENG. 1400 NORTH GOODMAN STREET ROCHESTER, N.Y. 14603-0450 TEL: 716-338-5474 FAX: 716-338-8959	MKT/GS REG AFF QA DCO NUMBER DATE
THE INFORMATION LISTED BELOW MUST APPEAR ON ALL VENDOR PROOFS (COLORKEYS, MATCHPRINTS/CHROMALINS) OR THEY WILL BE REJECTED			

PLACED PICTURES

AW #LB9135/06	DESCRIPTION Boston Lens II & IV Pat	ient Inst./Pkg. Insert	SPEC. NO. 4549
NUMBER OF PRINT		ER OF PMS COLORS 1	TOTAL AW PAGES 2
PMS COLORS		PROCESS MATCH COLO	ORS
1. PMS 326 AQUA			

PACKAGE INSERT

Boston® IV (itafocon B) Boston[®] II (itafocon A)

Spherical & Aspherical Contact Lenses for Myopia & Hyperopia, Keratoconus Contact Lenses. Bifocal Contact Lenses for Presbyopia and Toric Lenses to Correct Astigmatism in Not aphakic and Aphakic Persons

> Contact Lenses Daily Wear

IMPORTANT: this information for future use. This package insert is intended but should be made available to patients upon request. The provide the patient with the prescribed lens.

CAUTION: Federal (USA) Law restricts this device to sale by or on the order

DESCRIPTION	
Boston® IV (itafocon B) and Boston® are Rigid Gas Permeable Contact Lens composed of methyl-methacrylate-di conate-siloxanyl copolymer. Boston IV materials are available in clear and blu Boston IV lens materials contain the c D & C Green No. 6. Boston II lens mat available in clear, blue and green. The lens material contain the color additiv. No. 6; the green Boston II lens material color additive D & C Green No. 6 and Yellow No. 18.	materials, methylita- / lens le. The blue color additive erials are blue Boston re D & C Gree ial contain the
Boston IV & Boston II Contact Lenses spherical shell of the following dimen	

		Light manismiccance whole.	00111111 23070
Spherical Lens Design		Water Content	<1%
Power Range Diameter Base Curve Range	-20.00D to +20.00D in 0.25D increments 7.0 mm to 11.5 mm 5.00 mm to 9.00 mm in .01 mm increments	Oxygen Permeability Wetting Angle *gas to gas method **polarographic method	18* (11**) 20° (ISO/Fatt)
Aspherical Lens Designs (Some of these designs are patented; manufacture of these lenses in Boston IV (itafocon B) and/or Boston II (itafocon A) materials are authorized to licensed labs only)		Boston IV and Boston II C placed on the cornea act focus light rays on the ret vides a more even surfac	as a refracting medium to cina. The toric lens pro- e over the uneven astig-

Power Range Diameter Base Curve Range Bifocal Lens Designs (Some of these designs are patented;

Power Range Diameter Base Curve Range Segment Heights -2.00 mm to +1.00 mm Add Powers +1.00D to +3.75D Prism Ballast 0.5 to 3.5 prism diopte

Base Curve Range

Base Optic Zone

7.0 mm to 11.5 mm
4.00 mm to 9.00 mm
in 0.01 mm increments
5.00 mm to 9.00 mm
in 0.01 mm increments

4.01 contact lenses or using contact lens solution which is to be used to care for Boston IV or Boston IV contact lens materials.

4.02 mm to 11.5 mm
4.00 mm to 9.00 mm
in 0.01 mm increments

than Keratoconus, that affects the cornea, conjunctiva, or eyelids · Severe insufficiency of lacrimal secretion (dry if not aphakic Any systemic disease that may affect the eye or Allergic reactions of ocular surfaces or adnexa that may be induced or exaggerated by wearing

Boston IV

PHYSICAL PROPERTIES

Oxygen Permeability

PHYSICAL PROPERTIES

These lenses are indicated for daily wear. These

non-aphakic persons with non-diseased eyes that

are myopic, hyperopic or presbyopic, or have kera

astigmatism that does not interfere with visual acuity

The lenses may be disinfected using a chemical dis

CONTRAINDICATIONS (REASONS NOT TO USE)

DO NOT USE either Boston IV or Boston II Contact

Lenses when any of the following conditions exist:

Acute or subacute inflammation of the anterior

Any eye disease, injury, or abnormality, other

Specific Gravity

Refractive Index

Water Content

Specific Gravity

Refractive Index

the retina.

INDICATIONS (USES)

nfection system only.

Red or irritated eyes

WARNINGS Patients should be advised of the following warnings pertaining to contact lens wear:

• Problems with contact lenses and lens care products could result in serious injury to the eve. It practitioner's directions and all labeling instructions for proper use of lenses and lens care prodincluding corneal ulcers, can develop rapidly and

lead to loss of vision. Daily wear lenses are not indicated for overnight wear lenses while sleeping. Clinical studies have shown that the risk of serious adverse reaction worn overnight.

 Studies have shown that contact lens wearers who are smokers have a higher incidence of • If a patient experiences eye discomfort, excessiv tearing, vision changes, or redness of the eye, the patient should be instructed to immediately

eyecare practitioner. PRECAUTIONS Practitioner Note: Boston® IV and Boston® II contact lenses are not sterile when shipped from the Authorized Boston® Manufacturer. Prior to dis-

> ing to the appropriate lens care regimen. • Due to the small number of patients enrolled in clinical investigation of lenses, all refractive pow available in the lens material are not evaluated in significant numbers. Consequently, when cting an appropriate lens design and parameters, the eyecare practitioner should consider all characteristics of the lens that can affect lens

pensing, clean and disinfect the lens(es) accord

permeability, wettability, central and peripheral thickness, and optic zone diameter. • The potential impact of these factors on the patient's ocular health should be carefully weighed against the patient's need for refractive correction; therefore, the continuing ocular health of the patient and lens performance on the eye should be carefully monitored by the

• Patients who wear contact lenses to correct presbyonia may not achieve the best corrected visual acuity for either far or near vision. Visual requirements vary with the individual and should

• Aphakic patients should not be fitted with either the eye. Boston IV or Boston II Contact Lenses until the • Carefully follow the handling, insertion, removal. determination is made that the eye has healed completely. Before leaving the everage practitioner's office

tions in the Patient Instructions for Boston IV and the patient should be able to properly remove lenses or should have someone else available who s can remove the lenses for him or her.

. Eyecare practitioners should instruct the patien to remove the lenses immediately if the eve

patients about the following care regimen and

er, and not all solutions are safe for use with all

and/or lenses. Keep them away from extreme - Always use fresh unexpired lens care

solutions. for the use of contact lens solutions. - Use only a chemical (not heat) lens care system. Use

Boston IV or Boston II Contact Lenses. - Do not use saliva or anything other than the ting lenses.

- Sterile unpreserved solutions, when used. should be discarded after the time specified in the labeling directions. - Always keep the lenses completely immersed in

lenses are not being worn (stored). If dry storage is desired to store the lenses for a longer d of time, they must first be cleaned rinsed with water and carefully dried by blotting with a soft lint-free tissue prior to being these lenses should be cleaned and disinfected prior to insertion.

 If the lens sticks (stops moving) on the eye, follow the recommended directions on Care for a Sticking Lens. The lens should move freely on the eye for the continued health of the eye. If non-movement of the lens continues, the patient should be instructed to immediately consult his

· Always wash and rinse hands before handling lens es. Do not get cosmetics, lotions, soaps, creams, It is best to put on lenses before putting on make up. Water-based cosmetics are less likely to dam-

• Do not touch contact lenses with the fingers or hands if the hands are not free of foreign materioccur, causing distorted vision and/or injury to

cleaning, disinfecting, storing and wearing instructhe lenses and consult the evecare practitioner.

• If aerosol products such as hair spray are used while wearing lenses, exercise caution and keep eyes closed until the spray has settled. Always handle lenses gently and avoid dropping

 Avoid all harmful or irritating vapors and fumes while wearing lenses. Ask your evecare practitioner about wearing lens-

them on hard surfaces.

es during water activities and other sports. • Instruct patient to inform his or her health care practitioner (doctor) that the patient wears con-

 Never use tweezers or other tools to remove lenses from the lens case unless specifically indicated for that use. Pour the lens into the hand • Do not touch the lens with fingernails.

 Instruct the patient to contact his or her evecare practitioner before using any medicine in the Instruct the patient to inform his or her employe

that he or she wears contact lenses. Some jobs may require use of eye protection equipment or may require that the patient not wear contact · As with any contact lens, follow-up visits are necessary to assure the continuing health of the

as to a recommended follow-up schedule. ADVERSE EFFECTS The patient should be informed that the following problems may occur:

• Eyes stinging, burning, itching (irritation), or other eye pain Comfort is less than when lens was first placed

• Feeling of something in the eye such as a foreign body, scratched area Excessive watering (tearing) of the eyes

 Unusual eye secretions • Redness of the eyes Reduced sharpness of vision (poor visual acuity) Blurred vision, rainbows, or halos around objects Sensitivity to light (photophobia)

 Dry eyes If the patient notices any of the above, he or she Immediately remove lenses.

 If the discomfort or problem stops, then look
 Basic Instructions: Always wash, rinse, and dry hands before handling contact lenses. closely at the lens. If the lens is in any way damthe lens in the storage case and contact the eyecare practitioner. If the lens has dirt, an evelash, and the lens appears undamaged, the patient should thoroughly clean, rinse, and disinfect the lenses; then reinsert them. After reinsertion, if the problem continues, **immediately remove**

When any of the above problems occur, a serious condition such as infection, corneal ulcer, neovascularization, or iritis may be present. The patier should be instructed to keep the lens off the eve and seek immediate professional identifica

avoid serious eye damage. FITTING

extremely important.

apply to Boston® IV (itafocon B) and Boston® II tion of the fitting techniques, refer to Boston IV/ Boston II Professional Fitting and Information Guide

6 to 10 hours

enses are NOT intended for overnight (extended)

information and specific instructions on the lens

(First Clean and Rinse. Then Disinfect Lenses

Always use fresh unexpired lens care solutions.

chemical (not heat) and carefully follow instruc-

alternate or mix lens care systems unless

by the eyecare practitioner

indicated on solution labeling, or if advised

Use the recommended system of lens care,

care regimen recommended for the patient:

General Lens Care

ed by Bausch & Lomb for use with either Boston IV Practitioner Marketing Representative or Boston II Contact Lenses. Eyecare practitioners Boston Products Group of Bausch & Lomb may recommend alternate products that are appro-1400 North Goodman Street priate for the patient's use with his or her lens(es). LENS CARE TABLE 1-800-225-1241

Professional Fitting Guides are also available through your Authorized Boston® Manufacturer WEARING SCHEDULE Disinfect The wearing and replacement schedules Boston ADVANCE® Comfor should be determined by the eyecare practitioner. Patients tend to overwear the lenses initially Boston ADVANCE® Comfort importance of adhering to the initial maximum

wearing schedule. Regular checkups, as deter-Boston® Conditioning Solution nined by the eyecare practitioner, are also Boston IV and Boston II Contact Lenses are indicat ed for daily wear. The suggested wearing time for

Lubricate/Rewet Boston® Rewetting Drops Weekly Boston® ONE STEP Enzymatic Liquid Enzymatic Cleaner Cleaner . Note: Some solutions may have more than one

12 to All Waking Hours Read the label on the solution bottle, and follow *if the lenses continue to be well-tolerated WARNING: Boston IV and Boston II Contact

• Do not use saliva or anything other than the rec- Clean one lens first (always the same lens first) to avoid mix-ups), rinse the lens thoroughly lenses. Do not put lenses in the mouth. Lenses should be cleaned, rinsed, and disinfed remove the cleaning solution, mucus, and film

rinsing are necessary to r Then repeat the procedure for the second lens from the lens surface. Disinfecting is necessary to destroy harmful germs. After cleaning disinfect lenses using the sys-. Always remove, clean, rinse, and disinfect lenses

Lens Care System

according to the schedule prescribed by the eye-care practitioner. The use of an enzyme or a clear and/or the eyecare practitioner. Follow the instructions provided in the disinfecting soluing solution does not substitute for The lens care products listed below are recommen

• To store lenses, disinfect and leave them in the closed/unopened case until ready to wear. If lenses are not to be used immediately follow-ing disinfection, the patient should be instructed to consult the package insert or the eve-

 After removing the lenses from the lens case. empty and rinse the lens storage case with solution as recommended by the lens case manufacturer: then allow the lens case to air storage solution. Replace lens case at regular intervals as recommended by the lens case

cating/rewetting solution which can be used worn to make them more comfortable. · Evecare practitioners may recommend a to effectively remove protein deposits from Boston® IV or Boston® II Contact Lenses.

 Boston IV and Boston II Contact Lenses canno be heat (thermally) disinfected.

LENS CASE CLEANING AND MAINTENANCE

Contact lens cases can be a source of bacterial growth, Lens cases should be emptied, cleaned, from the lens surface, and put that lens into case manufacturer or the eyecare practitioner, and allowed to air dry. Lens cases should be replaced at regular intervals as recommended by the lens case manufacturer or the eyecare

practitioner. CARE FOR A STICKING (NONMOVING) LENS If the lens sticks (stops moving/cannot be

apply one to three drops of a recommended lubricating or rewetting solution directly to the eye and eye before removing it. If nonmovement of the lens continues after 5 minutes, the patient should **EMERGENCIES**

The patients should be informed that if chemicals of any kind (household products, gardening solu the eyes, the patient should: FLUSH EYES IMMEDI-ATELY WITH TAP WATER THEN REMOVE LENSES PROMPTLY, IF POSSIBLE, AND IMMEDIATELY CON-TACT THE EYECARE PRACTITIONER OR VISIT A HOSPI-TAL EMERGENCY ROOM WITHOUT DELAY.

HOW SUPPLIED

Each lens is supplied (nonsterile) in a plastic lens curve, diopter power, diameter, center thickness color, UV-absorber (if present) and lot number. height, prism ballast and truncation may be included for bifocal lenses.

REPORTING OF ADVERSE REACTIONS

wearing either Boston® IV or Boston® II Contact enses or adverse experiences with the lenses should be reported to: Bausch & Lomb 1400 North Goodman Street Rochester, NY 14609 1-800-333-4730

© Bausch & Lomb Incorporate Rochester, NY 14609 1-800-333-4730

Print Date: 09/07

BACK SIDE

THIS COMPONENT PRINTED TWO SIDES

APPROVALS / DATE PKG/GRPH	
T MKT/GS REG AFF QA DCO NUMBER	
DATE	
HR	HROMALINS) OR THEY WILL BE REJECT

PLACED PICTURES

AW #LB9135/06 DESCRIPTION Boston Lens II & IV	V Patient Inst./Pkg. Insert SPEC. NO. 4549
	IMBER OF PMS COLORS 1 TOTAL AW PAGES 2
PMS COLORS	PROCESS MATCH COLORS
1. PMS 326 AQUA	

PACKAGE INSERT

this information for future use. This package insert is intended but should be made available to patients upon request. The

CAUTION: Federal (USA) Law restricts this device to sale by or on the order

DESCRIPTION	
are Rigid Gas Permeal composed of methyl-conate-siloxanyl copo materials are available Boston IV Iens materia D & C Green No. 6. Bc available in clear, blue lens material contain I No. 6; the green Bost	B) and Boston® II (itafocon A ole Contact Lens materials, methacrylate-dimethylita- lymer. Boston IV lens in clear and blue. The blue als contain the color additive sston II lens materials are and green. The blue Boston the color additive D & C Gree on II lens material contain th reen No. 6 and C.I. Solvent
Boston IV & Boston II spherical shell of the t	Contact Lenses are hemi- following dimensions:
Spheric	cal Lens Design
Power Range	-20.00D to +20.00D in 0.25D increments

	:- 0.0FD :
Diameter	in 0.25D increments
Diameter	7.0 mm to 11.5 mm
Base Curve Range	5.00 mm to 9.00 mm
	in .01 mm increments
Aspherical	Lens Designs
(Some of these d	esigns are patented;
manufacture of the	se lenses in Boston IV
(itafocon B) and/or	Boston II (itafocon A)
materials are authorize	ed to licensed labs only)
Power Range	-20.00D to +20.00D
	in 0.25D increments
Diameter	7.0 mm to 11.5 mm
Base Curve Range	6.00 mm to 9.20 mm
base carve kange	in 0.01 mm increments

Bifocal Lens Designs (Some of these designs are patented; manufacture of these lenses in Boston IV (itafocon B) and/or Boston II (itafocon A) Power Range Diameter Base Curve Range Segment Heights Add Powers +1.00D to +3.75D

Prism Ballast 0.5 to 3.5 prism diopte Base Curve Range

7.0 mm to 11.5 mm
4.00 mm to 9.00 mm
in 0.01 mm increments
5.00 mm to 9.00 mm
in 0.01 mm increments

4.01 contact lenses or using contact lens solution which is to be used to care for Boston IV or Boston IV contact lens materials.

4.02 mm to 11.5 mm
4.00 mm to 9.00 mm
in 0.01 mm increments Base Optic Zone

> Patients should be advised of the following warnings pertaining to contact lens wear:

Boston IV PHYSICAL PROPERTIES Specific Gravity Refractive Index Water Content Oxygen Permeability

matic cornea and thus helps to focus light rays on

ing to the appropriate lens care regimen. • Due to the small number of patients enrolled in clinical investigation of lenses, all refractive pow available in the lens material are not evaluated in significant numbers. Consequently, when cting an appropriate lens design and parameters, the eyecare practitioner should consider all characteristics of the lens that can affect lens

PRECAUTIONS

permeability, wettability, central and peripheral thickness, and optic zone diameter. • The potential impact of these factors on the patient's ocular health should be carefully weighed against the patient's need for refractive correction; therefore, the continuing ocular health of the patient and lens performance on the eye should be carefully monitored by the

 Patients who wear contact lenses to correct presbyopia may not achieve the best corrected visual acuity for either far or near vision. Visual requirements vary with the individual and should

• Aphakic patients should not be fitted with either the eye. Boston IV or Boston II Contact Lenses until the • Carefully follow the handling, insertion, removal.

the patient should be able to properly remove lenses or should have someone else available who s

. Eyecare practitioners should instruct the patien to remove the lenses immediately if the eve

er, and not all solutions are safe for use with all - Do not heat the conditioning/storage solution

- Always use fresh unexpired lens care

for the use of contact lens solutions. - Use only a chemical (not heat) lens care system. Use

The patient should be informed that the following problems may occur: other eye pain

 Comfort is less than when lens was first placed • Feeling of something in the eye such as a foreign body, scratched area

 Unusual eye secretions • Redness of the eyes Reduced sharpness of vision (poor visual acuity) Blurred vision, rainbows, or halos around objects

Dry eyes

When any of the above problems occur, a serious condition such as infection, corneal ulcer, neovascularization, or iritis may be present. The patier should be instructed to keep the lens off the eve and seek immediate professional identifica

 Ask your evecare practitioner about wearing lenses during water activities and other sports.

• Instruct patient to inform his or her health care practitioner (doctor) that the patient wears con- Never use tweezers or other tools to remove lenses from the lens case unless specifically indi-

• If aerosol products such as hair spray are used

eyes closed until the spray has settled.

them on hard surfaces.

while wearing lenses.

while wearing lenses, exercise caution and keep

Always handle lenses gently and avoid dropping

Avoid all harmful or irritating vapors and fumes

cated for that use. Pour the lens into the hand • Do not touch the lens with fingernails. Instruct the patient to contact his or her evecare practitioner before using any medicine in the

 Instruct the patient to inform his or her employe that he or she wears contact lenses. Some jobs may require use of eye protection equipment or may require that the patient not wear contact

· As with any contact lens, follow-up visits are necessary to assure the continuing health of the as to a recommended follow-up schedule. ADVERSE EFFECTS

• Eyes stinging, burning, itching (irritation), or

Excessive watering (tearing) of the eyes

Sensitivity to light (photophobia)

If the patient notices any of the above, he or she Immediately remove lenses.

 If the discomfort or problem stops, then look
 Basic Instructions: closely at the lens. If the lens is in any way damthe lens in the storage case and contact the eyecare practitioner. If the lens has dirt, an evelash, and the lens appears undamaged, the patient should thoroughly clean, rinse, and disinfect the

avoid serious eye damage.

FITTING

apply to Boston® IV (itafocon B) and Boston® II tion of the fitting techniques, refer to Boston IV/ Boston II Professional Fitting and Information Guide copies of which are available from: Practitioner Marketing Representative

1-800-225-1241

Professional Fitting Guides are also available

The wearing and replacement schedules

should be determined by the eyecare practi-

tioner. Patients tend to overwear the lenses initially

importance of adhering to the initial maximum

wearing schedule. Regular checkups, as deter-

nined by the eyecare practitioner, are also

Boston IV and Boston II Contact Lenses are indicat

ed for daily wear. The suggested wearing time for

WEARING SCHEDULE

extremely important.

through your Authorized Boston® Manufacturer

Boston Products Group of Bausch & Lomb 1400 North Goodman Street priate for the patient's use with his or her lens(es). LENS CARE TABLE

> Disinfect Boston ADVANCE® Comfor Boston ADVANCE® Comfort

> > Lubricate/Rewet Boston® Rewetting Drops Weekly Boston® ONE STEP Enzymatic Liquid Enzymatic Cleaner Cleaner

. Note: Some solutions may have more than one Read the label on the solution bottle, and follow

ed to consult the package insert or the eve-

 After removing the lenses from the lens case. empty and rinse the lens storage case with solution as recommended by the lens case manufacturer: then allow the lens case to air storage solution. Replace lens case at regular intervals as recommended by the lens case • Eyecare practitioners may recommend a lubri

worn to make them more comfortable. · Evecare practitioners may recommend a Boston® IV or Boston® II Contact Lenses. LENS CASE CLEANING AND MAINTENANCE

Contact lens cases can be a source of bacterial growth, Lens cases should be emptied, cleaned, case manufacturer or the eyecare practitioner, and allowed to air dry. Lens cases should be replaced at regular intervals as recommended by the lens case manufacturer or the eyecare

practitioner. CARE FOR A STICKING (NONMOVING) LENS If the lens sticks (stops moving/cannot be

apply one to three drops of a recommended lubricating or rewetting solution directly to the eye and eye before removing it. If nonmovement of the lens continues after 5 minutes, the patient should **EMERGENCIES**

The patients should be informed that if chemicals of any kind (household products, gardening solu the eyes, the patient should: FLUSH EYES IMMEDI-ATELY WITH TAP WATER THEN REMOVE LENSES PROMPTLY, IF POSSIBLE, AND IMMEDIATELY CON-TACT THE EYECARE PRACTITIONER OR VISIT A HOSPI-TAL EMERGENCY ROOM WITHOUT DELAY.

HOW SUPPLIED

Each lens is supplied (nonsterile) in a plastic lens curve, diopter power, diameter, center thickness color, UV-absorber (if present) and lot number. height, prism ballast and truncation may be included for bifocal lenses.

REPORTING OF ADVERSE REACTIONS

wearing either Boston® IV or Boston® II Contact enses or adverse experiences with the lenses should be reported to: Bausch & Lomb 1400 North Goodman Street Rochester, NY 14609 1-800-333-4730

Rochester, NY 14609 1-800-333-4730

Boston® IV (itafocon B) Boston[®] II (itafocon A)

Spherical & Aspherical Contact Lenses for Myopia & Hyperopia, Keratoconus Contact Lenses. Bifocal Contact Lenses for Presbyopia and Toric Lenses to Correct Astigmatism in Not aphakic and Aphakic Persons

Contact Lenses Daily Wear

IMPORTANT: provide the patient with the prescribed lens.

ici ci i ici ita	ACTIONS		
11.5 mm 9.00 mm	*gas to gas method **polarographic method (ISO/Fatt)		
+20.00D rements	Wetting Angle	20°	
	Water Content Oxygen Permeability	<1% 18* (11**)	
re hemi- ions:	Light Transmittance (Above 500 nm)		
	Refractive Index	1.471	
	Specific Gravity	1.13	

-2.00 mm to +1.00 mm

 Red or irritated eyes WARNINGS

PHYSICAL PROPERTIES

Boston IV and Boston II Contact Lenses when placed on the cornea act as a refracting medium to focus light rays on the retina. The toric lens provides a more even surface over the uneven astig-

INDICATIONS (USES) These lenses are indicated for daily wear. These lenses are intended to correction of visual acuity in non-aphakic persons with non-diseased eyes that are myopic, hyperopic or presbyopic, or have kera

astigmatism that does not interfere with visual acuity The lenses may be disinfected using a chemical dis nfection system only. CONTRAINDICATIONS (REASONS NOT TO USE) DO NOT USE either Boston IV or Boston II Contact

Lenses when any of the following conditions exist: Acute or subacute inflammation of the anterior Any eye disease, injury, or abnormality, other than Keratoconus, that affects the cornea, conjunctiva, or eyelids · Severe insufficiency of lacrimal secretion (dry

if not aphakic Any systemic disease that may affect the eye or Allergic reactions of ocular surfaces or adnexa that may be induced or exaggerated by wearing

> determination is made that the eye has healed completely. Before leaving the everage practitioner's office can remove the lenses for him or her.

tions for proper use of lenses and lens care prodpatients about the following care regimen and

lead to loss of vision. Daily wear lenses are not indicated for overnight wear lenses while sleeping. Clinical studies have shown that the risk of serious adverse reaction and/or lenses. Keep them away from extreme

worn overnight. Studies have shown that contact lens wearers solutions. who are smokers have a higher incidence of

• Problems with contact lenses and lens care prod-

ucts could result in serious injury to the eve. It

practitioner's directions and all labeling instruc-

including corneal ulcers, can develop rapidly and

pensing, clean and disinfect the lens(es) accord

• If a patient experiences eye discomfort, excessive tearing, vision changes, or redness of the eye, the patient should be instructed to immediately Boston IV or Boston II Contact Lenses. eyecare practitioner.

- Do not use saliva or anything other than the ting lenses. Practitioner Note: Boston® IV and Boston® II - Sterile unpreserved solutions, when used. contact lenses are not sterile when shipped from the Authorized Boston® Manufacturer. Prior to disshould be discarded after the time specified in

the labeling directions. - Always keep the lenses completely immersed in lenses are not being worn (stored). If dry storage is desired to store the lenses for a longer d of time, they must first be cleaned rinsed with water and carefully dried by blotting with a soft lint-free tissue prior to being these lenses should be cleaned and disinfected

prior to insertion. If the lens sticks (stops moving) on the eye, follow the recommended directions on Care for a Sticking Lens. The lens should move freely on the eye for the continued health of the eye. If non-movement of the lens continues, the patient should be instructed to immediately consult his

· Always wash and rinse hands before handling lens es. Do not get cosmetics, lotions, soaps, creams, It is best to put on lenses before putting on make up. Water-based cosmetics are less likely to dam-

• Do not touch contact lenses with the fingers or hands if the hands are not free of foreign materioccur, causing distorted vision and/or injury to

cleaning, disinfecting, storing and wearing instruc-

tions in the Patient Instructions for Boston IV and

lenses; then reinsert them. After reinsertion, if the problem continues, **immediately remove** the lenses and consult the evecare practitioner. • Do not use saliva or anything other than the reclenses. Do not put lenses in the mouth. Lenses should be cleaned, rinsed, and disinfed

rinsing are necessary to r from the lens surface. Disinfecting is necessary to destroy harmful germs. . Always remove, clean, rinse, and disinfect lenses according to the schedule prescribed by the eye-care practitioner. The use of an enzyme or a clear

ing solution does not substitute for The lens care products listed below are recommen ed by Bausch & Lomb for use with either Boston IV or Boston II Contact Lenses. Eyecare practitioners may recommend alternate products that are appro-

Lens Care System

Boston® Conditioning Solution

*if the lenses continue to be well-tolerated WARNING: Boston IV and Boston II Contact enses are NOT intended for overnight (extended)

6 to 10 hours

12 to All Waking Hours

information and specific instructions on the lens care regimen recommended for the patient: General Lens Care (First Clean and Rinse. Then Disinfect Lenses

Always wash, rinse, and dry hands before handling contact lenses. Always use fresh unexpired lens care solutions. · Use the recommended system of lens care, chemical (not heat) and carefully follow instruc-

alternate or mix lens care systems unless

by the eyecare practitioner

indicated on solution labeling, or if advised

 Clean one lens first (always the same lens first) to avoid mix-ups), rinse the lens thoroughly remove the cleaning solution, mucus, and film from the lens surface, and put that lens into Then repeat the procedure for the second lens

 After cleaning disinfect lenses using the sysand/or the eyecare practitioner. Follow the instructions provided in the disinfecting solu-

• To store lenses, disinfect and leave them in the closed/unopened case until ready to wear. If lenses are not to be used immediately follow-ing disinfection, the patient should be instruct-

cating/rewetting solution which can be used

to effectively remove protein deposits from Boston IV and Boston II Contact Lenses canno be heat (thermally) disinfected.

© Bausch & Lomb Incorporate

Print Date: 09/07

BACK SIDE